

04 Health procedures

04.5 Poorly children

- If a child appears unwell during the day, for example has a raised temperature, sickness, diarrhoea or pains, particularly in the head or stomach then the setting manager calls the parents and asks them to collect the child or send a known carer to collect on their behalf.
- If a child has a raised temperature, they are kept cool by removing top clothing and kept away from draughts. Under parents consent, we will administer Calpol. Parents are called to collect their child.
- A child's temperature is taken and checked regularly, using a forehead thermometer.
- In an emergency an ambulance is called and the parents are informed.
- Parents are advised to seek medical advice before returning them to the setting; the setting
 can refuse admittance to children who have a raised temperature, sickness and diarrhoea
 or a contagious infection or disease.
- Where children have been prescribed antibiotics for an infectious illness or complaint, parents are asked to keep them at home for 48 hours.
- After diarrhoea or vomiting, parents are asked to keep children home for 48 hours following the last episode.
- Some activities such as sand and water play and self-serve snack will be suspended for the duration of any outbreak.
- The setting has information about excludable diseases and exclusion times.
- The setting manager has a list of notifiable diseases and contacts Public Health England (PHE) and Ofsted in the event of an outbreak.
- If staff suspect that a child who falls ill whilst in their care is suffering from a serious disease that may have been contracted abroad such as Ebola, immediate medical assessment is required. The setting manager or deputy calls NHS111 and informs parents.



HIV virus, like other viruses such as Hepatitis, (A, B and C), are spread through body fluids. Hygiene precautions for dealing with body fluids are the same for all children and adults.

- Single use vinyl gloves and aprons are worn when changing children's nappies, pants and clothing that are soiled with blood, urine, faeces or vomit.
- Protective rubber gloves are used for cleaning/sluicing clothing after changing.
- Soiled clothing is rinsed and bagged for parents to collect.
- Spills of blood, urine, faeces or vomit are cleared using mild disinfectant solution and mops; cloths used are disposed of with clinical waste.
- Tables and other furniture or toys affected by blood, urine, faeces or vomit are cleaned using a disinfectant.

Further guidance

Good Practice in Early Years Infection Control (Pre-school Learning Alliance 2009)

Medication Administration Record (Early Years Alliance 2019)

Guidance on infection control in schools and other childcare settings (Public Health Agency) https://www.publichealth.hscni.net/sites/default/files/Guidance on infection control in%20s chools poster.pdf